

# ***MIT Joint Program on the Science and Policy of Global Change***



## **Influence of Air Quality Model Resolution on Uncertainty Associated with Health Impacts**

*Tammy M. Thompson and Noelle E. Selin*

**Report No. 208  
December 2011**

The MIT Joint Program on the Science and Policy of Global Change is an organization for research, independent policy analysis, and public education in global environmental change. It seeks to provide leadership in understanding scientific, economic, and ecological aspects of this difficult issue, and combining them into policy assessments that serve the needs of ongoing national and international discussions. To this end, the Program brings together an interdisciplinary group from two established research centers at MIT: the Center for Global Change Science (CGCS) and the Center for Energy and Environmental Policy Research (CEEPR). These two centers bridge many key areas of the needed intellectual work, and additional essential areas are covered by other MIT departments, by collaboration with the Ecosystems Center of the Marine Biology Laboratory (MBL) at Woods Hole, and by short- and long-term visitors to the Program. The Program involves sponsorship and active participation by industry, government, and non-profit organizations.

To inform processes of policy development and implementation, climate change research needs to focus on improving the prediction of those variables that are most relevant to economic, social, and environmental effects. In turn, the greenhouse gas and atmospheric aerosol assumptions underlying climate analysis need to be related to the economic, technological, and political forces that drive emissions, and to the results of international agreements and mitigation. Further, assessments of possible societal and ecosystem impacts, and analysis of mitigation strategies, need to be based on realistic evaluation of the uncertainties of climate science.

This report is one of a series intended to communicate research results and improve public understanding of climate issues, thereby contributing to informed debate about the climate issue, the uncertainties, and the economic and social implications of policy alternatives. Titles in the Report Series to date are listed on the inside back cover.


Ronald G. Prinn and John M. Reilly  
*Program Co-Directors*

For more information, please contact the Joint Program Office

Postal Address: Joint Program on the Science and Policy of Global Change  
77 Massachusetts Avenue  
MIT E19-411  
Cambridge MA 02139-4307 (USA)

Location: 400 Main Street, Cambridge  
Building E19, Room 411  
Massachusetts Institute of Technology

Access: Phone: +1(617) 253-7492  
Fax: +1(617) 253-9845  
E-mail: [globalchange@mit.edu](mailto:globalchange@mit.edu)  
Web site: <http://globalchange.mit.edu/>

 Printed on recycled paper

# Influence of Air Quality Model Resolution on Uncertainty Associated with Health Impacts

Tammy M. Thompson<sup>\*†</sup> and Noelle E. Selin<sup>\*</sup>

## Abstract

*We evaluate the uncertainty associated with regional air quality modeling grid resolution when calculating the health benefits of proposed air quality regulations. Using a regional photochemical model (CAMx), we ran two modeling episodes (a 2006 basecase and a 2018 attainment demonstration, both for Houston, Texas) at 36, 12, 4 and 2 km resolution. The basecase model performance was evaluated for each resolution for both monitor-based and population-weighted calculations of daily maximum 8-hour averaged ozone. Results from each resolution were more similar to each other than they are to actual measured values. However, the model performance improved when population weighted ozone concentration was used as the metric versus the standard daily maximum ozone concentrations at monitor site locations. Then population-weighted ozone concentrations were used to calculate the estimated health impacts of modeled ozone reduction from the basecase to the attainment demonstration including the 95% confidence intervals associated with each impact from concentration-response functions. We found that estimated avoided mortalities were not significantly different using coarse resolution, although 36 km resolution may over predict some potential health impacts. Given the cost/benefit analyses requirements of the Clean Air Act, the uncertainty associated with human health impacts and therefore the results reported in this study, we conclude that population weighted ozone concentrations obtained using regional photochemical models at 36 km resolution are meaningful relative to values obtained using fine (12 km or finer) resolution modeling. This result opens up the possibility for uncertainty analyses on 36 km resolution air quality modeling results, which are on average 10 times more computationally efficient.*

## Contents

1. INTRODUCTION.....	1
2. METHODS .....	4
2.1 Comprehensive Air Quality Model with Extensions (CAMx) .....	4
2.2 Variable Grid Performance Analysis.....	5
2.3 Health Impacts.....	6
3. COMPARISON OF MONITOR-BASED AND POPULATION-BASED PERFORMANCE EVALUATION.....	6
3.1 2006 Base Case: Monitor-based Analysis .....	6
3.2 2006 Base Case: Population-Weighted Analysis.....	7
3.3 2006 Basecase vs 2018 Control Case: Population-Weighted Comparison.....	8
4. UNCERTAINTY ANALYSIS OF HEALTH IMPACTS AT VARYING MODEL RESOLUTION .....	10
5. ERROR ANALYSIS AND APPLICABILITY TO OTHER REGIONS .....	11
6. CONCLUSIONS AND IMPLICATIONS FOR REGULATORY APPROACH .....	11
7. REFERENCES.....	12

## 1. INTRODUCTION

Ground level ozone air pollution has been linked to adverse human health impacts (EPA 2010a) and is regulated in the United States by the Environmental Protection Agency (EPA) with the goal of protecting health. In implementing these regulations, states use results from air quality models to demonstrate compliance (called attainment demonstrations). While many

---

<sup>\*</sup> MIT Joint Program for the Science and Policy of Global Change, Cambridge, MA.

<sup>†</sup> Corresponding Author, Email: [TammyT@mit.edu](mailto:TammyT@mit.edu)

elements of ozone concentration and impacts are uncertain (including emissions, chemistry, and health impacts), model resolution requirements make uncertainty analyses in attainment demonstrations unfeasible. However, understanding the true projected benefits of ozone regulation, as the Clean Air Act also mandates, requires uncertainty analysis. Here, we compare the variation associated with simulated ozone at various model resolutions with uncertainty in estimated human health impacts, applying population-weighted concentrations. We use the results of this analysis to evaluate the potential for using coarser-scale model resolution for uncertainty analyses of prospective ozone regulations.

Air quality modeling is applied extensively in the U.S. to assess implementation of air quality standards. Many U.S. counties are in non-attainment of the 2008 EPA National Ambient Air Quality Standards (NAAQS) standard of 75 ppb. Many more will be designated non-attainment if the standard is decreased in 2011, as planned. Non-attainment means that the three year average of the fourth highest 8-hour average ozone concentration (called a “design value”) at any monitor is greater than the standard. Where non-attainment occurs, states are charged with developing an attainment demonstration to show how they plan to meet the standard. Because the formation of ozone occurs in complex, non-linear chemical reactions between nitrogen oxides (NO<sub>x</sub>) and volatile organic compounds (VOCs), attainment demonstrations require air quality modeling to help develop and evaluate pollution control strategies. At a minimum, EPA requires a resolution of 12 km by 12 km or smaller for attainment demonstrations using approved air quality models (with coarse resolution grid cells extending over all potentially contributing sources), but recommends that each case be evaluated independently to identify the potential model prediction improvements associated with finer scale resolution.

Extensive analyses in the atmospheric chemistry literature have evaluated the impact of model resolution on ozone production (Arunachalam *et al.*, 2006; Cohen *et al.*, 2006; Jang *et al.*, 1995; Tie *et al.*, 2010; Valari & Menut, 2008). Eulerian photochemical air quality models instantly and homogeneously disperse low level emissions (including ozone precursors NO<sub>x</sub> and VOCs) throughout the grid cell. This spatial averaging impacts the chemistry by smoothing concentration gradients of precursors over large areas, which in some cases has shown to reduce modeled ozone titration effects and ozone formation hotspots. As a result, many studies have found that larger scale resolution (> 12 km grid cells) leads to an under-prediction of daily maximum 8 hour ozone averages, and an over-prediction of daily minimum 8 hour ozone averages (Jang *et al.*, 1995; Arunachalam *et al.*, 2006; Tie *et al.*, 2010). Some studies indicate that 12 km resolution is often not fine enough to capture sharp ozone concentration gradients that can occur near large sources of precursors, like power plants or dense urban areas with a lot of traffic (Valeri & Menut, 2008; Kumar & Russell, 1995).

While previous studies have assessed the errors in predicted ozone versus monitor concentrations, it is unclear what the impact of resolution-based errors are on predicted benefits of regulation. Section 812 of the Clean Air Act (CAA) requires the EPA to conduct a cost/benefit analysis of the impact of that act (EPA, 2011). As part of the latest periodic assessment, EPA conducted an uncertainty analysis on the CAA and the latest standards by comparing estimated

human health impacts of ozone and particulate matter concentrations in 2020 under the environmental regulation mandated by the CAA, versus likely concentrations of those two pollutants if the CAA were not implemented. While the uncertainty analysis addressed relative potential impacts of many uncertainties, probability distributions were included only for concentration response functions. With respect to atmospheric uncertainties, the EPA argued that errors in ozone benefits using a 12 km grid are likely minor, but primarily because the benefits are far outweighed by PM reductions; they did not quantitatively compare resolution. These types of analyses are not assessed during attainment demonstrations; previously, Wesson *et al.* (2010) noted this disconnect and argue that multi-pollutant strategies are more effective than traditional attainment strategies at reducing population risk. The National Research Council (NRC) has also criticized the lack of comprehensive uncertainty analyses in the evaluation of environmental policy options, calling for probabilistic multi-source uncertainty analyses (NRC, 2002).


A growing literature has used regional modeling to assess the potential impacts of global climate change and future emissions on ozone concentrations (Selin *et al.*, 2009; Chang *et al.*, 2010; Bell *et al.*, 2007; West *et al.*, 2007; Tagaris *et al.*, 2009; Knowlton *et al.* 2004). Bell *et al.* (2007) calculated a 0.11% to 0.27% increase (the 95% confidence interval) in percent change in mortality across 31 cities in the U.S. based on the difference between modeled maximum daily ozone concentrations in five summers each around 2050 and the 1990s. Similarly, Knowlton *et al.* (2004) projected a 4.5% increase on average in acute mortality in New York State in 2050 due to climate change. Tagaris *et al.* (2009) evaluated the uncertainty associated with meteorological conditions based on the range of temperature and humidity values modeled by several global change models, concluding that uncertainty due to meteorology was larger than uncertainty associated with human health impacts. All of these studies used resolution of 36 km or coarser from climate models. Given the stochastic nature of the climate system, ensemble analyses of future projected ozone changes, incorporating uncertainties, would be beneficial to produce probabilistic information for decision-making (NRC, 2002).

Here, we address the disconnect between methodologies for attainment demonstrations and for impacts analysis. We evaluate the impact on modeled potential ozone exposure and calculated human health response uncertainty resulting from the temporal and spatial smoothing seen in coarse grid domains (Jang *et al.*, 1995; Arunachalam *et al.*, 2006; Tie *et al.*, 2010) due to the spatial smoothing of ozone precursors, which can eliminate NO<sub>x</sub> titration and hot spot formation. We focus on comparing the relative uncertainty associated with model resolution and resulting predicted ambient concentrations, with uncertainty associated with projected human health impacts. We use this comparison to identify a resolution appropriate for impacts analysis uncertainty, taking into account relative errors and computational limitations.

## 2. METHODS

### 2.1 Comprehensive Air Quality Model with Extensions (CAMx)

We use CAMx ([www.camx.com](http://www.camx.com)), an EPA-approved regional air quality model (EPA, 2007). We use a well-documented air quality episode developed in part during the Texas Air Quality Study II (TexAQSI) (TCEQ, 2006; TCEQ, 2010b). The episode was created for the Houston/Galveston/Brazoria (HGB) non-attainment area and includes a 2006 basecase and a 2018 attainment demonstration scenario. Emissions inventories were speciated, and spatially and temporally processed using the Emissions Preprocessing System (EPS3). The 2006 basecase inventory represents actual 2006 emissions, while the 2018 emissions inventories were created to demonstrate HGB attainment with the 1997 ozone standard and includes proposed controls on ozone precursors (TCEQ, 2010a). Resolution of the original episode includes a coarse parent grid at 36 km, and three nested grids at 12 km, 4 km, and 2 km (**Figure 1**). Meteorological inputs are consistent in both scenarios and were developed using the fifth generation Penn State/NCAR mesoscale model MM5 (Grell *et al.*, 1994) for August 13-September 15, 2006; for the 2 km domain, meteorological data is interpolated by CAMx from 4 km. A detailed description of the episode is provided by the Texas Commission on Environmental Quality TCEQ (2010a). Performance of the episode was evaluated previously (TCEQ, 2010a), and met EPA criteria (+/- 15% for bias and error as described below for 1 hour ozone concentrations with a 60 ppb threshold).



**Figure 1.** CAMx Modeling domain. For each resolution only the modeling results within the area covered by the 2 km domain (the HGB area), shown above, are used.

## 2.2 Variable Grid Performance Analysis

We focus here on the HGB area (Figure 1, red square). We conduct four simulations each for the two cases (2006 basecase and 2018 control case), with increasingly coarse resolution over the HGB area (2 km, 4 km, 12 km, and 36 km). We evaluate the performance of the 2006 basecase in reproducing daily maximum 8 hour averaged ozone concentrations at air quality monitors in the region. This metric is selected for evaluation because it determines attainment and it is necessary for input into concentration-response functions for impact analysis; this metric is not required by EPA performance evaluation. We use the statistical measures Mean Normalized Bias (MNB) and Mean Normalized Gross Error (MNGE) as shown in Equation 1 and Equation 2 respectively.

$$MNB = \frac{1}{N} \sum_1^N \left( \frac{(Model - Obs)}{Obs} \right) * 100 \% \quad (1)$$

$$MNGE = \frac{1}{N} \sum_1^N \left( \frac{|Model - Obs|}{Obs} \right) * 100 \% \quad (2)$$

### 2.3 Health Impacts

For our analysis of health impacts and potential benefits, we use maximum daily population weighted concentrations ( $M_{Pop}$ ) as a surrogate for exposure for both model and measurement calculated using Equation 3.

$$M_{Pop} = \frac{\sum_g p_g \max \{c_g\}}{\sum_g p_g} \quad (3)$$


where  $p_g$  is the population in grid cell  $g$ , and  $c_g$  is the maximum 8 hour ozone concentration in grid cell  $g$ . Population distribution is from U.S. Census data, mapped using Geographical Information System (GIS) software. For the base case, year 2007 is used, and for the 2018 policy case, projected 2015 population is applied. For population-weighted analysis for monitor data, only those grid cells with monitors located in them are used in the calculation. This metric represents a rough but best available and commonly-used estimate for the potential for human exposure. In reality, exposure depends not only on the ambient concentration of pollutants at any given time and location, but also on the daily patterns of people being exposed: when, where and how they travel to and from activities and their initial health (EPA, 2010b). The potential impacts on human health of changes in ozone concentrations are calculated by multiplying population-weighted concentrations by concentration response functions for acute mortality, respiratory hospital admissions (adults over 65 yrs), respiratory symptom days, minor restricted activity day, asthma attacks, and bronchodilator usage. We use the survey of Bickel and Friedrich (2005) to specify these functions and related 95% confidence intervals, assuming a linear relationship between daily maximum 8 hour ozone concentrations and impacts, and no minimum health impact threshold. For baseline mortality rate, we used the average rate for developed countries from Lopez *et al.* (2006)

## 3. COMPARISON OF MONITOR-BASED AND POPULATION-BASED PERFORMANCE EVALUATION

### 3.1 2006 Basecase: Monitor-based Analysis

We first evaluated the performance of the 2006 basecase episode on the daily maximum 8 hour ozone concentrations modeled for each of the air quality monitors located in the HGB domain for each of the four spatial resolution runs. **Figure 2a** shows the error (Equation 2) in comparing monitor values to simulated concentrations in the grid cell containing the monitor (at each resolution). Error increases from 25% to 74% as model resolution increases from 2 km to 36 km. **Figure 2b** compares the difference between the three coarser resolutions relative to the 2 km fine scale modeling result; the difference in predicted ozone between coarser and finer scale resolution ranges from 1%-15%. We conclude from this comparison that results from the different resolutions are more similar to each other than they are to actual measured values.






**Figure 2. (a):** Model Data vs. Measured Data: Model error comparing CAMx results for four grid resolution runs for the 2006 basecase to measured concentrations at all air quality monitor sites in the 2 km domain. Bias and error results are approximately the same due to a high bias of the model, so only error is shown in Figure 2a. **(b):** "Coarse" Model Data vs. "Fine" Model Data: Model Error for modeled "coarse" resolution ozone concentrations compared with modeled 2k "fine" resolution ozone concentrations.

### 3.2 2006 Base Case: Population-weighted Analysis

To assess a metric more relevant to health impacts, we compared the ability of different model resolution to reproduce population-weighted concentration. **Figure 3** shows the impact of resolution on the population-weighted concentrations as modeled using the 2006 basecase. These results are compared to the measured concentrations at the monitors within the HGB domain. Higher-resolution modeling (4 or 2 km) exhibits no clear benefit in comparison with 12 km resolution when considering population-weighted concentrations. The 36 km simulation is biased high (by 3 ppb resolution relative to finer scale model results); however, on average

across all monitor locations, modeled concentrations are 10 ppb higher than measured concentrations.




**Figure 3.** Population-weighted maximum ozone concentration for each resolution from the 2006 episode compared to the population-weighted maximum ozone concentration calculated from the measured values at the monitors using the 2 km resolution.

The 2006 basecase episode over-estimates the population weighted ozone concentrations on most days (Figure 3). This bias is consistent with the monitor-based results presented in Section 3.1 above. Additionally, the model is not able to consistently capture the daily variability of the measured results. However, the results are improved over the standard performance evaluation statistics for 12 km and 36 km resolution as presented in Figure 2. The mean normalized error of the population weighted daily maximum 8 hour ozone concentrations modeled using the 2006 basecase (and only cells containing monitors) compared to population weighted measured concentrations at air quality monitors average across the episode is 26%, 27%, 24% and 32% for 2 km, 4 km, 12 km, and 36 km resolution respectively.


### 3.3 2006 Basecase vs 2018 Control Case: Population-weighted Comparison

We compared population-weighted ozone changes between the 2006 basecase with the 2018 control case, to identify the variation in concentration between different resolutions for benefits analysis. **Figure 4** shows the maximum population-weighted concentration (Equation 3) calculated for each day of both the 2006 and 2018 episodes, using the fine resolution (2 km) modeling runs. Also shown are the population-weighted concentrations using measured monitor data and the HGB domain.



**Figure 4.** Population weighted daily maximum 8 hour averaged ozone concentrations using fine scale 2 km population and concentration data from monitors, and modeled values from each of the runs.

**Figure 5** shows the change in population weighted 8 hour ozone concentrations from base case 2006 model data to control case 2018 model data. Based on these results, the control scenarios as proposed by the TCEQ in the 2018 episode clearly impact the modeled ozone concentrations, with an average 10 ppb decrease in both population weighted concentrations and maximum daily 8 hour ozone concentration from 2006 basecase results. The calculated ozone decrease differs depending on what model resolution is used: the average decrease is 8 ppb for both the 2 km and 4 km model resolutions, 10 ppb for the 12 km model resolution, and 12 ppb for the 36 km resolution. For comparison, the average change in the fourth highest daily maximum 8 hour ozone at all monitor locations only is 8 ppb, 7 ppb, 7 ppb and 6 ppb respectively.



**Figure 5.** Impact of 2018 control strategy (2018 Control case – 2006 Base case) on daily maximum 8 hour ozone population weighted ozone concentration by resolution.

#### 4. UNCERTAINTY ANALYSIS OF HEALTH IMPACTS AT VARYING MODEL RESOLUTION

We use the change in population weighted daily maximum 8 hour ozone shown above in Figure 5, averaged across all days of the episode, to calculate the expected health benefits from the Houston attainment demonstration. We use these values to compare the estimated benefits that would be calculated based on concentrations predicted at the four resolutions, and concentration-response functions and 95% confidence intervals as described above.

**Table 1** shows the calculated change in mortalities and morbidities between 2006-2018, based on concentration information from the four different modeling resolutions. Also shown are the total number of cases in 2006 calculated using data from air quality monitors. The mean (5) and 95% confidence interval (2-7) for the change in mortalities is identical up to 12 km. For 36 km resolution, the mean is 7 and 95% confidence interval 2-9. Basecase (2006) mortality is calculated to be between 15 and 58 deaths per ozone month (May-September) due to acute exposure, with a mean at 44.

For avoided respiratory hospital admissions (adults >65 years of age) and avoided bronchodilator usage, health benefits estimated using the 36 km resolution ozone modeling results fully contain the 95% confidence interval calculated using finer scale modeling results. However, for changes in respiratory symptom days, minor restricted activity days and asthma attacks, the 36 km 95% confidence interval does not fully contain the confidence intervals for the finer scale resolution results. For these endpoints, an analysis done at 36 km resolution could potentially over-estimate the benefits due to modeled ozone reductions. However, the EPA has

found that most of the monetary benefits associated with health improvements comes from reduction in mortalities due to the high value of a statistical life (EPA, 2011).

**Table 1.** Change in human health impacts due to the control scenarios proposed as part of the 2018 Houston Attainment Demonstration. The top row shows baseline human health impacts due to ozone measured at monitors in 2006.

Mean with 95% Confidence Interval	Change in Mortality (# of deaths in area)	Respiratory Hospital Admissions Adults >65 yrs	Respiratory Symptom Day	Minor Restricted Activity Day	Asthma Attack	Bronchodilator usage
Values Calculated Using Population Weighted Concentrations as Measured by Air Quality Monitors in 2006						
06 Monitor Data	44 (15,58)	207 (-83,497)	546442 (94385,1043208)	190427 (72859,314618)	71037 (5464,137438)	1208796 (-430530,2649416)
Change (Decrease) in Metrics between the 2006 Modeled Basecase and the 2018 Modeled Control Case						
Model 2k	5 (2,7)	25 (-10,60)	65466 (11308,124980)	22814 (8729,37692)	8511 (655,16466)	144818 (-51579,317409)
Model 4k	5 (2,7)	24 (-10,59)	64601 (11158,123330)	22513 (8613,37195)	8398 (646,16248)	142906 (-50898,313218)
Model 12k	5 (2,7)	23 (-9,56)	61302 (10589,117031)	21363 (8174,35295)	7969 (613,15418)	135607 (-48299,297222)
Model 36k	7 (2,9)	32 (-13,76)	83552 (14432,159508)	29117 (11140,48106)	10862 (836,21015)	184827 (-65829,405101)

## 5. ERROR ANALYSIS AND APPLICABILITY TO OTHER REGIONS

As a first test of whether our results may be applicable to other regions, we use the CAMx Process Analysis (PA) tool to calculate individual contributions from each physical and chemical process within the model, to the final concentration of ozone. This allows us to better understand the cause of the resolution-dependent differences in our study. We used PA and the python based Process Analysis (pyPA) post-processing tool, developed by Henderson *et al.* (2011) to analyze results for September 12<sup>th</sup> because that day had the largest difference between the 2006 basecase and the 2018 control case (Figure 5). Our results indicate that the resolution difference is due to chemistry: ozone production due to chemistry in the fine resolution model is twice that of the coarse resolution model during the hours of 10AM to 12PM (12 ppb per hour vs. 6 ppb per hour). As chemistry processes are consistent across different regions, this suggests that our analysis may be valid beyond the Houston area; however, further studies could address other regions as episodes are developed.

## 6. CONCLUSIONS AND IMPLICATIONS FOR REGULATORY APPROACH

To evaluate the uncertainty associated with air quality modeling resolution for calculating health benefits of proposed regulations, we ran two modeling episodes (a basecase and an attainment demonstration, both for Houston, Texas) at 36, 12, 4 and 2 km resolution. We evaluated basecase model performance for each resolution for both monitor-based and population-weighted calculations of 8-hour maximum ozone. We then used population-weighted ozone concentrations to calculate the estimated health impacts of ozone reduction from the basecase to the attainment demonstration including the 95% confidence intervals associated with each impact from concentration-response functions. We found that estimated avoided mortalities

were not significantly different using coarse resolution, although 3 km resolution may over predict some potential health impacts.

We evaluated the performance of the 2006 basecase with respect to monitor-based and population-weighted 8-hour maximum ozone. Results from each resolution were more similar to each other than they are to actual measured values. However, the model performance improved when population weighted ozone concentration was used as the metric versus the standard daily maximum ozone concentrations.

We compared the difference in the population weighted ozone concentrations between resolutions and between the 2006 base case and the 2018 control case. The coarse scale resolution (36 km) showed the largest decrease from basecase to attainment demonstration case. The average change in daily maximum 8 hour ozone population weighted concentrations are 10 ppb, 7 ppb, 8 ppb and 8 ppb for 36 km, 12 km, 4 km, and 2 km resolution respectively.

We used the population-weighted ozone concentration difference to calculate acute mortality, respiratory hospital admissions (adults over 65 yrs), respiratory symptom days, minor restricted activity day, asthma attacks, and bronchodilator usage. The confidence interval of the mortality predicted by the coarse scale resolution completely contains the confidence interval of the mortality predicted by the fine scale results.

Given the cost/benefit analyses requirements of the Clean Air Act, the uncertainty associated with human health impacts and therefore the results reported in Table 1, we conclude that population weighted ozone concentrations obtained using regional photochemical models at 36 km resolution are meaningful relative to values obtained using fine (12 km or finer) resolution modeling. This result opens up the possibility for uncertainty analyses on 36 km resolution air quality modeling results, which are on average 10 times more computationally efficient.

## **Acknowledgements**

The research described has been supported by the U.S. Environmental Protection Agency's STAR program through grant R834279 and by the MIT Joint Program on the Science and Policy of Global Change. It has not been subjected to any EPA review and therefore does not necessarily reflect the views of the Agency, and no official endorsement should be inferred.

## **7. REFERENCES**

- Arunachalam, S; A. Holland; B. Do; M. Abraczinskas. 2006: A quantitative assessment of the influence of grid resolution on predictions of future-year air quality in North Carolina, USA. *Atmos. Env.* 40(2006) 5010-5026.
- Bell, M.L.; R. Goldberg; C. Hogrefe; P.L. Kinney; K. Knowlton; B. Lynn; J. Rosenthal; C. Rosenzweig; J.A. Patz. 2007: Climate Change, Ambient Ozone, and Health in 50 US Cities. *Climate Change* **2007**, 1, 61-76.
- Bickel, P.; R. Friedrich (ed) 2005: ExternE: Externalities of Energy. Methodology 2005 Update.

- Chang, H.H.; J. Zhou; M. Fuentes. 2010: Impact of climate change on ambient ozone level and mortality in southeastern United States. *Int. J. Environ. Res. Public Health*. 7 (2010) 2866-2880.
- Cohen, D; Y. Hu; A. Russell. 2006: Dependence of ozone sensitivity analysis on grid resolution. *Atmos. Env.* 40 (2006) 126-135.
- Environ. 2010: Comprehensive Air Quality Model with Extensions User's Guide. Environ International Corp. March 2010.
- EPA. 2007: Guidance on the Use of Models and Other Analyses for Demonstrating Attainment of Air Quality Goals for Ozone, PM<sub>2.5</sub>, and Regional Haze. Office of Air Quality Planning and Standards. Report B-07-002 April 2007.
- EPA. 2010a: Ground-Level Ozone: Health and Environment. Office of Air and Radiation. <http://www.epa.gov/glo/health.html>.
- EPA. 2010b: Stochastic Human Exposure and Dose Simulation (SHEDS) Multimedia Model Version 3. Office of Research and Development: Human Exposure and Atmospheric Sciences. Accessed 2011. [http://www.epa.gov/heasd/products/sheds\\_multimedia/sheds\\_mm.html](http://www.epa.gov/heasd/products/sheds_multimedia/sheds_mm.html)
- EPA. 2011: The Benefits and Costs of the Clean Air Act from 1990 to 2020. Office of Air and Radiation. Second Prospective Study. Accessed May 2011. <http://www.epa.gov/oar/sect812/prospective2.html>
- Grell, G.; J. Dudhia; D.R. Stauffer. A description of the fifth generation Penn State/NCAR mesoscale model (MM5); Tech. Note NCAR/TN-398+STR; National Center for Atmospheric Research: Boulder, CO, 1994.
- Henderson, B.H.; Y. Kimura; E. McDonald-Buller; D.T. Allen; W. Vizuetta. 2011: Comparison of Lagrangian Process Analysis tools for Eulerian air quality models. *Atmos. Env.* 45 (2011) 5200-5211.
- Jang, J.C.; H.E. Jeffries; S. Tonnesen. 1995: Sensitivity of Ozone to Model Grid Resolution – II. Detailed Process Analysis for Ozone Chemistry. *Atmos. Env.* 29(21).
- Knowlton, K.; J.E. Rosenthal; C. Hogrefe; B. Lynn; S. Gaffin; R. Goldberg; C. Rosenzweig; J.Y. Ku; P.L. Kinney. 2004: Assessing Ozone-Related Health Impacts Under a Changing Climate. *Environ. Health Perspect.* 2004, 112, 1557-1563.
- Kumar, N; A. Russel. 1995: Multiscale air quality modeling of the Northeastern United States. *Atmos. Env.* 30(7) 1099-1116.
- Lopez, A.D.; C.D. Mathers; M. Ezzati; D.T. Jamison; C.J.L. Murray (ed). 2006: Global Burden of Disease and Risk Factors. New York: Oxford University Press.
- NRC (National Research Council). 2002: Estimating the Public Health Benefits of Proposed Air Pollution Regulations. Washington, DC: National Academies Press.
- Selin, N.E.; S. Wu; K.M. Nam; J.M. Reilly; S. Paltsev; R.G. Prinn; M.D. Webster. 2009: Global Health and economic impacts of future ozone pollution. *Environment Research Letters*. Vol, 4 (2009).
- Tagaris, E.; K.J. Liao; A.J. Delucia; L. Deck; P. Amar; A.G. Russell. 2009: Potential Impact of Climate Change on Air Pollution-Related Human Health Effects. *Environ. Sci. Technol.* 2009, 43, 4979-4988.
- TCEQ. 2006: TexAQSI Field Study. <http://www.tceq.texas.gov/airquality/research/texaqs>

- TCEQ. 2010a: Houston-Galveston-Brazoria Attainment Demonstration and Reasonable Further Progress State Implementation Plan Revisions for the 1997 Eight-Hour Ozone Standard. March 10, 2010. [http://www.tceq.texas.gov/airquality/sip/HGB\\_eight\\_hour.html](http://www.tceq.texas.gov/airquality/sip/HGB_eight_hour.html)
- TCEQ. 2010b: Air Quality Modeling Files ftp Site. Accessed January 2010. <ftp://ftp.tceq.state.tx.us/pub/OEPAA/TAD/Modeling/>
- Tie, X; G. Brasseur; Z. Ying. 2010: Impact of model resolution on chemical ozone formation in Mexico City: application of the WRF-Chem model. *Atmos. Chem. Phys.* 10, 8983-8995.
- Valari, M; L. Menut. 2008: Does an increase in air quality models' resolution bring surface ozone concentrations closer to reality? *J. Atmos. Oceanic. Tech.* (25) 1955-1968.
- Wesson, K.; N. Fann; M. Morris; T. Fox; B. Hubbell. 2010: A multi-pollutant, risk-based approach to air quality management: Case study for Detroit. *Atmospheric Pollution Research* 1 (2010) 296-304. <http://www.atmospolres.com/articles/Volume1/issue4/APR-10-037.pdf>
- West, J.J.; S. Szopa; D.A. Hauglustaine. 2007: Human Mortality Effects of Future Concentrations of Tropospheric Ozone. *C. R. Geosci.* 2007, 339, 775-783.



## REPORT SERIES of the MIT *Joint Program on the Science and Policy of Global Change*

1. **Uncertainty in Climate Change Policy Analysis**  
*Jacoby & Prinn* December 1994
2. **Description and Validation of the MIT Version of the GISS 2D Model** *Sokolov & Stone* June 1995
3. **Responses of Primary Production and Carbon Storage to Changes in Climate and Atmospheric CO<sub>2</sub> Concentration** *Xiao et al.* October 1995
4. **Application of the Probabilistic Collocation Method for an Uncertainty Analysis** *Webster et al.* January 1996
5. **World Energy Consumption and CO<sub>2</sub> Emissions: 1950-2050** *Schmalensee et al.* April 1996
6. **The MIT Emission Prediction and Policy Analysis (EPPA) Model** *Yang et al.* May 1996 (*superseded* by No. 125)
7. **Integrated Global System Model for Climate Policy Analysis** *Prinn et al.* June 1996 (*superseded* by No. 124)
8. **Relative Roles of Changes in CO<sub>2</sub> and Climate to Equilibrium Responses of Net Primary Production and Carbon Storage** *Xiao et al.* June 1996
9. **CO<sub>2</sub> Emissions Limits: Economic Adjustments and the Distribution of Burdens** *Jacoby et al.* July 1997
10. **Modeling the Emissions of N<sub>2</sub>O and CH<sub>4</sub> from the Terrestrial Biosphere to the Atmosphere** *Liu* Aug. 1996
11. **Global Warming Projections: Sensitivity to Deep Ocean Mixing** *Sokolov & Stone* September 1996
12. **Net Primary Production of Ecosystems in China and its Equilibrium Responses to Climate Changes**  
*Xiao et al.* November 1996
13. **Greenhouse Policy Architectures and Institutions**  
*Schmalensee* November 1996
14. **What Does Stabilizing Greenhouse Gas Concentrations Mean?** *Jacoby et al.* November 1996
15. **Economic Assessment of CO<sub>2</sub> Capture and Disposal**  
*Eckaus et al.* December 1996
16. **What Drives Deforestation in the Brazilian Amazon?**  
*Pfaff* December 1996
17. **A Flexible Climate Model For Use In Integrated Assessments** *Sokolov & Stone* March 1997
18. **Transient Climate Change and Potential Croplands of the World in the 21st Century** *Xiao et al.* May 1997
19. **Joint Implementation: Lessons from Title IV's Voluntary Compliance Programs** *Atkeson* June 1997
20. **Parameterization of Urban Subgrid Scale Processes in Global Atm. Chemistry Models** *Calbo et al.* July 1997
21. **Needed: A Realistic Strategy for Global Warming**  
*Jacoby, Prinn & Schmalensee* August 1997
22. **Same Science, Differing Policies; The Saga of Global Climate Change** *Skolnikoff* August 1997
23. **Uncertainty in the Oceanic Heat and Carbon Uptake and their Impact on Climate Projections**  
*Sokolov et al.* September 1997
24. **A Global Interactive Chemistry and Climate Model**  
*Wang, Prinn & Sokolov* September 1997
25. **Interactions Among Emissions, Atmospheric Chemistry & Climate Change** *Wang & Prinn* Sept. 1997
26. **Necessary Conditions for Stabilization Agreements**  
*Yang & Jacoby* October 1997
27. **Annex I Differentiation Proposals: Implications for Welfare, Equity and Policy** *Reiner & Jacoby* Oct. 1997
28. **Transient Climate Change and Net Ecosystem Production of the Terrestrial Biosphere**  
*Xiao et al.* November 1997
29. **Analysis of CO<sub>2</sub> Emissions from Fossil Fuel in Korea: 1961-1994** *Choi* November 1997
30. **Uncertainty in Future Carbon Emissions: A Preliminary Exploration** *Webster* November 1997
31. **Beyond Emissions Paths: Rethinking the Climate Impacts of Emissions Protocols** *Webster & Reiner* November 1997
32. **Kyoto's Unfinished Business** *Jacoby et al.* June 1998
33. **Economic Development and the Structure of the Demand for Commercial Energy** *Judson et al.* April 1998
34. **Combined Effects of Anthropogenic Emissions and Resultant Climatic Changes on Atmospheric OH**  
*Wang & Prinn* April 1998
35. **Impact of Emissions, Chemistry, and Climate on Atmospheric Carbon Monoxide** *Wang & Prinn* April 1998
36. **Integrated Global System Model for Climate Policy Assessment: Feedbacks and Sensitivity Studies**  
*Prinn et al.* June 1998
37. **Quantifying the Uncertainty in Climate Predictions**  
*Webster & Sokolov* July 1998
38. **Sequential Climate Decisions Under Uncertainty: An Integrated Framework** *Valverde et al.* September 1998
39. **Uncertainty in Atmospheric CO<sub>2</sub> (Ocean Carbon Cycle Model Analysis)** *Holian* Oct. 1998 (*superseded* by No. 80)
40. **Analysis of Post-Kyoto CO<sub>2</sub> Emissions Trading Using Marginal Abatement Curves** *Ellerman & Decaux* Oct. 1998
41. **The Effects on Developing Countries of the Kyoto Protocol and CO<sub>2</sub> Emissions Trading**  
*Ellerman et al.* November 1998
42. **Obstacles to Global CO<sub>2</sub> Trading: A Familiar Problem**  
*Ellerman* November 1998
43. **The Uses and Misuses of Technology Development as a Component of Climate Policy** *Jacoby* November 1998
44. **Primary Aluminum Production: Climate Policy, Emissions and Costs** *Harnisch et al.* December 1998
45. **Multi-Gas Assessment of the Kyoto Protocol**  
*Reilly et al.* January 1999
46. **From Science to Policy: The Science-Related Politics of Climate Change Policy in the U.S.** *Skolnikoff* January 1999
47. **Constraining Uncertainties in Climate Models Using Climate Change Detection Techniques**  
*Forest et al.* April 1999
48. **Adjusting to Policy Expectations in Climate Change Modeling** *Shackley et al.* May 1999
49. **Toward a Useful Architecture for Climate Change Negotiations** *Jacoby et al.* May 1999
50. **A Study of the Effects of Natural Fertility, Weather and Productive Inputs in Chinese Agriculture**  
*Eckaus & Tso* July 1999
51. **Japanese Nuclear Power and the Kyoto Agreement**  
*Babiker, Reilly & Ellerman* August 1999
52. **Interactive Chemistry and Climate Models in Global Change Studies** *Wang & Prinn* September 1999

Contact the Joint Program Office to request a copy. The Report Series is distributed at no charge.

REPORT SERIES of the MIT *Joint Program on the Science and Policy of Global Change*

53. **Developing Country Effects of Kyoto-Type Emissions Restrictions** Babiker & Jacoby October 1999
54. **Model Estimates of the Mass Balance of the Greenland and Antarctic Ice Sheets** Bugnion Oct 1999
55. **Changes in Sea-Level Associated with Modifications of Ice Sheets over 21st Century** Bugnion October 1999
56. **The Kyoto Protocol and Developing Countries** Babiker et al. October 1999
57. **Can EPA Regulate Greenhouse Gases Before the Senate Ratifies the Kyoto Protocol?** Bugnion & Reiner November 1999
58. **Multiple Gas Control Under the Kyoto Agreement** Reilly, Mayer & Harnisch March 2000
59. **Supplementarity: An Invitation for Monopsony?** Ellerman & Sue Wing April 2000
60. **A Coupled Atmosphere-Ocean Model of Intermediate Complexity** Kamenkovich et al. May 2000
61. **Effects of Differentiating Climate Policy by Sector: A U.S. Example** Babiker et al. May 2000
62. **Constraining Climate Model Properties Using Optimal Fingerprint Detection Methods** Forest et al. May 2000
63. **Linking Local Air Pollution to Global Chemistry and Climate** Mayer et al. June 2000
64. **The Effects of Changing Consumption Patterns on the Costs of Emission Restrictions** Lahiri et al. Aug 2000
65. **Rethinking the Kyoto Emissions Targets** Babiker & Eckaus August 2000
66. **Fair Trade and Harmonization of Climate Change Policies in Europe** Viguier September 2000
67. **The Curious Role of "Learning" in Climate Policy: Should We Wait for More Data?** Webster October 2000
68. **How to Think About Human Influence on Climate** Forest, Stone & Jacoby October 2000
69. **Tradable Permits for Greenhouse Gas Emissions: A primer with reference to Europe** Ellerman Nov 2000
70. **Carbon Emissions and The Kyoto Commitment in the European Union** Viguier et al. February 2001
71. **The MIT Emissions Prediction and Policy Analysis Model: Revisions, Sensitivities and Results** Babiker et al. February 2001 (*superseded* by No. 125)
72. **Cap and Trade Policies in the Presence of Monopoly and Distortionary Taxation** Fullerton & Metcalf March '01
73. **Uncertainty Analysis of Global Climate Change Projections** Webster et al. Mar. '01 (*superseded* by No. 95)
74. **The Welfare Costs of Hybrid Carbon Policies in the European Union** Babiker et al. June 2001
75. **Feedbacks Affecting the Response of the Thermohaline Circulation to Increasing CO<sub>2</sub>** Kamenkovich et al. July 2001
76. **CO<sub>2</sub> Abatement by Multi-fueled Electric Utilities: An Analysis Based on Japanese Data** Ellerman & Tsukada July 2001
77. **Comparing Greenhouse Gases** Reilly et al. July 2001
78. **Quantifying Uncertainties in Climate System Properties using Recent Climate Observations** Forest et al. July 2001
79. **Uncertainty in Emissions Projections for Climate Models** Webster et al. August 2001
80. **Uncertainty in Atmospheric CO<sub>2</sub> Predictions from a Global Ocean Carbon Cycle Model** Holian et al. September 2001
81. **A Comparison of the Behavior of AO GCMs in Transient Climate Change Experiments** Sokolov et al. December 2001
82. **The Evolution of a Climate Regime: Kyoto to Marrakech** Babiker, Jacoby & Reiner February 2002
83. **The "Safety Valve" and Climate Policy** Jacoby & Ellerman February 2002
84. **A Modeling Study on the Climate Impacts of Black Carbon Aerosols** Wang March 2002
85. **Tax Distortions and Global Climate Policy** Babiker et al. May 2002
86. **Incentive-based Approaches for Mitigating Greenhouse Gas Emissions: Issues and Prospects for India** Gupta June 2002
87. **Deep-Ocean Heat Uptake in an Ocean GCM with Idealized Geometry** Huang, Stone & Hill September 2002
88. **The Deep-Ocean Heat Uptake in Transient Climate Change** Huang et al. September 2002
89. **Representing Energy Technologies in Top-down Economic Models using Bottom-up Information** McFarland et al. October 2002
90. **Ozone Effects on Net Primary Production and Carbon Sequestration in the U.S. Using a Biogeochemistry Model** Felzer et al. November 2002
91. **Exclusionary Manipulation of Carbon Permit Markets: A Laboratory Test** Carlén November 2002
92. **An Issue of Permanence: Assessing the Effectiveness of Temporary Carbon Storage** Herzog et al. December 2002
93. **Is International Emissions Trading Always Beneficial?** Babiker et al. December 2002
94. **Modeling Non-CO<sub>2</sub> Greenhouse Gas Abatement** Hyman et al. December 2002
95. **Uncertainty Analysis of Climate Change and Policy Response** Webster et al. December 2002
96. **Market Power in International Carbon Emissions Trading: A Laboratory Test** Carlén January 2003
97. **Emissions Trading to Reduce Greenhouse Gas Emissions in the United States: The McCain-Lieberman Proposal** Paltsev et al. June 2003
98. **Russia's Role in the Kyoto Protocol** Bernard et al. Jun '03
99. **Thermohaline Circulation Stability: A Box Model Study** Lucarini & Stone June 2003
100. **Absolute vs. Intensity-Based Emissions Caps** Ellerman & Sue Wing July 2003
101. **Technology Detail in a Multi-Sector CGE Model: Transport Under Climate Policy** Schafer & Jacoby July 2003
102. **Induced Technical Change and the Cost of Climate Policy** Sue Wing September 2003
103. **Past and Future Effects of Ozone on Net Primary Production and Carbon Sequestration Using a Global Biogeochemical Model** Felzer et al. (revised) January 2004

Contact the Joint Program Office to request a copy. The Report Series is distributed at no charge.

## REPORT SERIES of the MIT *Joint Program on the Science and Policy of Global Change*

104. **A Modeling Analysis of Methane Exchanges Between Alaskan Ecosystems and the Atmosphere** *Zhuang et al.* November 2003
105. **Analysis of Strategies of Companies under Carbon Constraint** *Hashimoto* January 2004
106. **Climate Prediction: *The Limits of Ocean Models*** *Stone* February 2004
107. **Informing Climate Policy Given Incommensurable Benefits Estimates** *Jacoby* February 2004
108. **Methane Fluxes Between Terrestrial Ecosystems and the Atmosphere at High Latitudes During the Past Century** *Zhuang et al.* March 2004
109. **Sensitivity of Climate to Diapycnal Diffusivity in the Ocean** *Dalan et al.* May 2004
110. **Stabilization and Global Climate Policy** *Sarofim et al.* July 2004
111. **Technology and Technical Change in the MIT EPPA Model** *Jacoby et al.* July 2004
112. **The Cost of Kyoto Protocol Targets: *The Case of Japan*** *Paltsev et al.* July 2004
113. **Economic Benefits of Air Pollution Regulation in the USA: *An Integrated Approach*** *Yang et al.* (revised) Jan. 2005
114. **The Role of Non-CO<sub>2</sub> Greenhouse Gases in Climate Policy: *Analysis Using the MIT IGSM*** *Reilly et al.* Aug. '04
115. **Future U.S. Energy Security Concerns** *Deutch* Sep. '04
116. **Explaining Long-Run Changes in the Energy Intensity of the U.S. Economy** *Sue Wing* Sept. 2004
117. **Modeling the Transport Sector: *The Role of Existing Fuel Taxes in Climate Policy*** *Paltsev et al.* November 2004
118. **Effects of Air Pollution Control on Climate** *Prinn et al.* January 2005
119. **Does Model Sensitivity to Changes in CO<sub>2</sub> Provide a Measure of Sensitivity to the Forcing of Different Nature?** *Sokolov* March 2005
120. **What Should the Government Do To Encourage Technical Change in the Energy Sector?** *Deutch* May '05
121. **Climate Change Taxes and Energy Efficiency in Japan** *Kasahara et al.* May 2005
122. **A 3D Ocean-Seaice-Carbon Cycle Model and its Coupling to a 2D Atmospheric Model: *Uses in Climate Change Studies*** *Dutkiewicz et al.* (revised) November 2005
123. **Simulating the Spatial Distribution of Population and Emissions to 2100** *Asadoorian* May 2005
124. **MIT Integrated Global System Model (IGSM) Version 2: *Model Description and Baseline Evaluation*** *Sokolov et al.* July 2005
125. **The MIT Emissions Prediction and Policy Analysis (EPPA) Model: *Version 4*** *Paltsev et al.* August 2005
126. **Estimated PDFs of Climate System Properties Including Natural and Anthropogenic Forcings** *Forest et al.* September 2005
127. **An Analysis of the European Emission Trading Scheme** *Reilly & Paltsev* October 2005
128. **Evaluating the Use of Ocean Models of Different Complexity in Climate Change Studies** *Sokolov et al.* November 2005
129. **Future Carbon Regulations and Current Investments in Alternative Coal-Fired Power Plant Designs** *Sekar et al.* December 2005
130. **Absolute vs. Intensity Limits for CO<sub>2</sub> Emission Control: *Performance Under Uncertainty*** *Sue Wing et al.* January 2006
131. **The Economic Impacts of Climate Change: *Evidence from Agricultural Profits and Random Fluctuations in Weather*** *Deschenes & Greenstone* January 2006
132. **The Value of Emissions Trading** *Webster et al.* Feb. 2006
133. **Estimating Probability Distributions from Complex Models with Bifurcations: *The Case of Ocean Circulation Collapse*** *Webster et al.* March 2006
134. **Directed Technical Change and Climate Policy** *Otto et al.* April 2006
135. **Modeling Climate Feedbacks to Energy Demand: *The Case of China*** *Asadoorian et al.* June 2006
136. **Bringing Transportation into a Cap-and-Trade Regime** *Ellerman, Jacoby & Zimmerman* June 2006
137. **Unemployment Effects of Climate Policy** *Babiker & Eckaus* July 2006
138. **Energy Conservation in the United States: *Understanding its Role in Climate Policy*** *Metcalfe* Aug. '06
139. **Directed Technical Change and the Adoption of CO<sub>2</sub> Abatement Technology: *The Case of CO<sub>2</sub> Capture and Storage*** *Otto & Reilly* August 2006
140. **The Allocation of European Union Allowances: *Lessons, Unifying Themes and General Principles*** *Buchner et al.* October 2006
141. **Over-Allocation or Abatement? *A preliminary analysis of the EU ETS based on the 2006 emissions data*** *Ellerman & Buchner* December 2006
142. **Federal Tax Policy Towards Energy** *Metcalfe* Jan. 2007
143. **Technical Change, Investment and Energy Intensity** *Kratena* March 2007
144. **Heavier Crude, Changing Demand for Petroleum Fuels, Regional Climate Policy, and the Location of Upgrading Capacity** *Reilly et al.* April 2007
145. **Biomass Energy and Competition for Land** *Reilly & Paltsev* April 2007
146. **Assessment of U.S. Cap-and-Trade Proposals** *Paltsev et al.* April 2007
147. **A Global Land System Framework for Integrated Climate-Change Assessments** *Schlosser et al.* May 2007
148. **Relative Roles of Climate Sensitivity and Forcing in Defining the Ocean Circulation Response to Climate Change** *Scott et al.* May 2007
149. **Global Economic Effects of Changes in Crops, Pasture, and Forests due to Changing Climate, CO<sub>2</sub> and Ozone** *Reilly et al.* May 2007
150. **U.S. GHG Cap-and-Trade Proposals: *Application of a Forward-Looking Computable General Equilibrium Model*** *Gurgel et al.* June 2007
151. **Consequences of Considering Carbon/Nitrogen Interactions on the Feedbacks between Climate and the Terrestrial Carbon Cycle** *Sokolov et al.* June 2007

Contact the Joint Program Office to request a copy. The Report Series is distributed at no charge.

## REPORT SERIES of the MIT *Joint Program on the Science and Policy of Global Change*

- 152. Energy Scenarios for East Asia: 2005-2025** *Paltsev & Reilly* July 2007
- 153. Climate Change, Mortality, and Adaptation: Evidence from Annual Fluctuations in Weather in the U.S.** *Deschênes & Greenstone* August 2007
- 154. Modeling the Prospects for Hydrogen Powered Transportation Through 2100** *Sandoval et al.* February 2008
- 155. Potential Land Use Implications of a Global Biofuels Industry** *Gurgel et al.* March 2008
- 156. Estimating the Economic Cost of Sea-Level Rise** *Sugiyama et al.* April 2008
- 157. Constraining Climate Model Parameters from Observed 20<sup>th</sup> Century Changes** *Forest et al.* April 2008
- 158. Analysis of the Coal Sector under Carbon Constraints** *McFarland et al.* April 2008
- 159. Impact of Sulfur and Carbonaceous Emissions from International Shipping on Aerosol Distributions and Direct Radiative Forcing** *Wang & Kim* April 2008
- 160. Analysis of U.S. Greenhouse Gas Tax Proposals** *Metcalf et al.* April 2008
- 161. A Forward Looking Version of the MIT Emissions Prediction and Policy Analysis (EPPA) Model** *Babiker et al.* May 2008
- 162. The European Carbon Market in Action: Lessons from the first trading period** Interim Report *Convery, Ellerman, & de Perthuis* June 2008
- 163. The Influence on Climate Change of Differing Scenarios for Future Development Analyzed Using the MIT Integrated Global System Model** *Prinn et al.* September 2008
- 164. Marginal Abatement Costs and Marginal Welfare Costs for Greenhouse Gas Emissions Reductions: Results from the EPPA Model** *Holak et al.* November 2008
- 165. Uncertainty in Greenhouse Emissions and Costs of Atmospheric Stabilization** *Webster et al.* November 2008
- 166. Sensitivity of Climate Change Projections to Uncertainties in the Estimates of Observed Changes in Deep-Ocean Heat Content** *Sokolov et al.* November 2008
- 167. Sharing the Burden of GHG Reductions** *Jacoby et al.* November 2008
- 168. Unintended Environmental Consequences of a Global Biofuels Program** *Melillo et al.* January 2009
- 169. Probabilistic Forecast for 21<sup>st</sup> Century Climate Based on Uncertainties in Emissions (without Policy) and Climate Parameters** *Sokolov et al.* January 2009
- 170. The EU's Emissions Trading Scheme: A Proto-type Global System?** *Ellerman* February 2009
- 171. Designing a U.S. Market for CO<sub>2</sub>** *Parsons et al.* February 2009
- 172. Prospects for Plug-in Hybrid Electric Vehicles in the United States & Japan: A General Equilibrium Analysis** *Karplus et al.* April 2009
- 173. The Cost of Climate Policy in the United States** *Paltsev et al.* April 2009
- 174. A Semi-Empirical Representation of the Temporal Variation of Total Greenhouse Gas Levels Expressed as Equivalent Levels of Carbon Dioxide** *Huang et al.* June 2009
- 175. Potential Climatic Impacts and Reliability of Very Large Scale Wind Farms** *Wang & Prinn* June 2009
- 176. Biofuels, Climate Policy and the European Vehicle Fleet** *Gitiaux et al.* August 2009
- 177. Global Health and Economic Impacts of Future Ozone Pollution** *Selin et al.* August 2009
- 178. Measuring Welfare Loss Caused by Air Pollution in Europe: A CGE Analysis** *Nam et al.* August 2009
- 179. Assessing Evapotranspiration Estimates from the Global Soil Wetness Project Phase 2 (GSWP-2) Simulations** *Schlosser and Gao* September 2009
- 180. Analysis of Climate Policy Targets under Uncertainty** *Webster et al.* September 2009
- 181. Development of a Fast and Detailed Model of Urban-Scale Chemical and Physical Processing** *Cohen & Prinn* October 2009
- 182. Distributional Impacts of a U.S. Greenhouse Gas Policy: A General Equilibrium Analysis of Carbon Pricing** *Rausch et al.* November 2009
- 183. Canada's Bitumen Industry Under CO<sub>2</sub> Constraints** *Chan et al.* January 2010
- 184. Will Border Carbon Adjustments Work?** *Winchester et al.* February 2010
- 185. Distributional Implications of Alternative U.S. Greenhouse Gas Control Measures** *Rausch et al.* June 2010
- 186. The Future of U.S. Natural Gas Production, Use, and Trade** *Paltsev et al.* June 2010
- 187. Combining a Renewable Portfolio Standard with a Cap-and-Trade Policy: A General Equilibrium Analysis** *Morris et al.* July 2010
- 188. On the Correlation between Forcing and Climate Sensitivity** *Sokolov* August 2010
- 189. Modeling the Global Water Resource System in an Integrated Assessment Modeling Framework: IGSM-WRS** *Strzepek et al.* September 2010
- 190. Climatology and Trends in the Forcing of the Stratospheric Zonal-Mean Flow** *Monier and Weare* January 2011
- 191. Climatology and Trends in the Forcing of the Stratospheric Ozone Transport** *Monier and Weare* January 2011
- 192. The Impact of Border Carbon Adjustments under Alternative Producer Responses** *Winchester* February 2011
- 193. What to Expect from Sectoral Trading: A U.S.-China Example** *Gavard et al.* February 2011
- 194. General Equilibrium, Electricity Generation Technologies and the Cost of Carbon Abatement** *Lanz and Rausch* February 2011
- 195. A Method for Calculating Reference Evapotranspiration on Daily Time Scales** *Farmer et al.* February 2011

Contact the Joint Program Office to request a copy. The Report Series is distributed at no charge.

**REPORT SERIES of the MIT *Joint Program on the Science and Policy of Global Change***

- 196. Health Damages from Air Pollution in China** *Matus et al.* March 2011
- 197. The Prospects for Coal-to-Liquid Conversion: A General Equilibrium Analysis** *Chen et al.* May 2011
- 198. The Impact of Climate Policy on U.S. Aviation** *Winchester et al.* May 2011
- 199. Future Yield Growth: What Evidence from Historical Data** *Gitiaux et al.* May 2011
- 200. A Strategy for a Global Observing System for Verification of National Greenhouse Gas Emissions** *Prinn et al.* June 2011
- 201. Russia's Natural Gas Export Potential up to 2050** *Paltsev* July 2011
- 202. Distributional Impacts of Carbon Pricing: A General Equilibrium Approach with Micro-Data for Households** *Rausch et al.* July 2011
- 203. Global Aerosol Health Impacts: Quantifying Uncertainties** *Selin et al.* August 2011
- 204. Implementation of a Cloud Radiative Adjustment Method to Change the Climate Sensitivity of CAM3** *Sokolov and Monier* September 2011
- 205. Quantifying the Likelihood of Regional Climate Change: A Hybridized Approach** *Schlosser et al.* Oct 2011
- 206. Process Modeling of Global Soil Nitrous Oxide Emissions** *Saikawa et al.* October 2011
- 207. The Influence of Shale Gas on U.S. Energy and Environmental Policy** *Jacoby et al.* November 2011
- 208. Influence of Air Quality Model Resolution on Uncertainty Associated with Health Impacts** *Thompson and Selin* December 2011