

What GHG Concentration Targets are Reachable in this Century?

Sergey Paltsev, John Reilly and Andrei Sokolov



Report No. 247
July 2013

The MIT Joint Program on the Science and Policy of Global Change is an organization for research, independent policy analysis, and public education in global environmental change. It seeks to provide leadership in understanding scientific, economic, and ecological aspects of this difficult issue, and combining them into policy assessments that serve the needs of ongoing national and international discussions. To this end, the Program brings together an interdisciplinary group from two established research centers at MIT: the Center for Global Change Science (CGCS) and the Center for Energy and Environmental Policy Research (CEEPR). These two centers bridge many key areas of the needed intellectual work, and additional essential areas are covered by other MIT departments, by collaboration with the Ecosystems Center of the Marine Biology Laboratory (MBL) at Woods Hole, and by short- and long-term visitors to the Program. The Program involves sponsorship and active participation by industry, government, and non-profit organizations.


To inform processes of policy development and implementation, climate change research needs to focus on improving the prediction of those variables that are most relevant to economic, social, and environmental effects. In turn, the greenhouse gas and atmospheric aerosol assumptions underlying climate analysis need to be related to the economic, technological, and political forces that drive emissions, and to the results of international agreements and mitigation. Further, assessments of possible societal and ecosystem impacts, and analysis of mitigation strategies, need to be based on realistic evaluation of the uncertainties of climate science.

This report is one of a series intended to communicate research results and improve public understanding of climate issues, thereby contributing to informed debate about the climate issue, the uncertainties, and the economic and social implications of policy alternatives. Titles in the Report Series to date are listed on the inside back cover.

Ronald G. Prinn and John M. Reilly
Program Co-Directors

For more information, please contact the Joint Program Office

Postal Address: Joint Program on the Science and Policy of Global Change
77 Massachusetts Avenue
MIT E19-411
Cambridge MA 02139-4307 (USA)
Location: 400 Main Street, Cambridge
Building E19, Room 411
Massachusetts Institute of Technology
Access: Phone: +1.617. 253.7492
Fax: +1.617.253.9845
E-mail: globalchange@mit.edu
Web site: <http://globalchange.mit.edu/>

 Printed on recycled paper

What GHG Concentration Targets are Reachable in this Century?

Sergey Paltsev^{*†}, John Reilly^{*}, and Andrei Sokolov^{*}

Abstract

We offer simulations that help to understand the relationship between GHG emissions and concentrations, and the relative role of long-lived (e.g., CO₂) and short-lived (e.g., CH₄) emissions. We show that, absent technologies to remove CO₂ from the atmosphere, the 350 CO₂ ppm target is out of reach in this century, even if all emissions drop to zero almost immediately (i.e. in 2015). A 350 ppm CO₂-equivalent target is potentially achievable, but would require CH₄ concentrations falling below preindustrial levels, and thus elimination of emissions from human activities such as rice and livestock agriculture. More realistically, even some of the most aggressive targets proposed through 2035 would lead to concentrations (CO₂ or CO₂-eq) in the 415–450 ppm range. This is only feasible if after 2035 emissions continued a downward path toward zero. Only in these cases would the temperature target of no more than 2 °C above preindustrial be achieved, and only after peaking above that level before declining.

Contents

1. INTRODUCTION.....	1
2. EMISSION SCENARIOS	2
3. RESULTING ATMOSPHERIC CONCENTRATIONS.....	3
4. RADIATIVE FORCING AND TEMPERATURE EFFECTS.....	6
5. CONCLUSIONS	8
6. REFERENCES.....	8

1. INTRODUCTION

Given the growing evidence that the world faces substantial risks associated with changing climate there are increasing questions about what concentrations of atmospheric carbon dioxide and greenhouse gases are achievable. Some have called for an atmospheric target for CO₂ as low as 350 ppm (Hansen et al., 2008). Since CO₂ levels are already reaching 400 ppm this means concentrations would need to fall. Some authors note the general difficulty people have in grasping the relationship between changes in flow (emissions) and stock (concentrations) as it applies to the climate problem (Sterman, 2008).

In this paper we provide several scenarios to illustrate the link between GHG emission trajectories and the targets proposed by the international community to address the challenges of climate change. The United Nations Framework Convention on Climate Change (UNFCCC) has reached an accord in 2009 for the so called Copenhagen pledges (United Nations, 2009), which are further specified in the Cancun agreements (United Nations, 2010). Even though, the targets in these agreements are provided mostly for 2020, they also express a longer-term objective of keeping the average global temperature rise below two degrees Celsius relative to the preindustrial levels.

^{*} Joint Program on the Science and Policy of Global Change, Massachusetts Institute of Technology, Cambridge, MA, USA.

[†] Corresponding author (email: paltsev@mit.edu).

This paper is organized in the following way. In the next section we discuss the scenarios that we consider to illustrate emissions trajectories and the proposed targets. Section 3 provides a discussion about the resulting atmospheric concentrations. In Section 4 we discuss the radiative forcing and temperature effects. Section 5 concludes.

2. EMISSION SCENARIOS

There are several studies that look at the temperature impacts of eliminating emissions (e.g., Matthews and Caldeira, 2008; Solomon et al., 2009; Gillett et al., 2011; Matthews and Zickfeld, 2012). Most of the studies focus on CO₂ emissions, neglecting or highly simplifying the effects of non-CO₂ greenhouse gases and aerosols. Here we use the MIT Integrated Global Systems Model (IGSM) that tracks GHG gases and aerosols. The MIT IGSM (Sokolov et al., 2005; Prinn et al., 2011) is an integrated assessment model (IAM) model of intermediate complexity, i.e. it represents the dynamics of the atmosphere and ocean in less detail than conventional general circulation models (GCMs), but goes beyond the approach taken by atmospheric energy balance models (EBMs) or ocean box models by using sophisticated parameterizations of the unresolved flows or by explicitly resolving the equations of geophysical fluid dynamics, albeit at coarse spatial resolution.

The scenarios considered in this study were motivated by a proposed abatement scenario (de Frondeville, 2012) through 2035 that was put forward as a candidate scenario for achieving a concentration level of 350 ppm of CO₂. We consider five emissions scenarios (**Figure 1**) using the MIT IGSM. These include: (1) a Copenhagen Scenario reflecting commitments under the UNFCCC through 2020 and extended through 2100 (Paltsev et al., 2012); (2) the Copenhagen Scenario with zero anthropogenic emissions after 2035 (Natural Only after 2035); (3) the Copenhagen Scenario with zero anthropogenic emissions immediately (Natural Only after 2015); (4) the IEA 450 ppm CO₂ Scenario with other gases constrained proportionally to CO₂ through the International Energy Agency's (IEA) horizon of 2035 (IEA, 2011) and then dropping linearly to zero at the rate for the decade 2025–2035; and (5) an Alternative Abatement Scenario proposed to us as an alternative to the IEA 450 Scenario intended to achieve 350 ppm of atmospheric CO₂ concentrations. In this latter scenario, cumulative emissions from 2010 to 2035 in the alternative scenario are about 77% of cumulative emissions in the IEA 450 Scenario, and the number 350 is about 77% of 450.

The heuristic used here to relate emissions and concentrations appears to be that a reduction in cumulative emissions would give a proportional reduction in concentrations. We make no attempt to evaluate the economic or political feasibility of these scenarios. Those that drop emissions to zero in one year are purposely extreme and intended to illustrate the earth system response to such a change. We have not evaluated the economic implications of the IEA 450 Scenario and the Alternative Abatement Scenario but these were developed with the idea that, at least through 2035, there were technological options that could lead to these outcomes. We contrast these to the internationally agreed goals in the Copenhagen Scenario and see that those commitments are far from adequate to achieve even the IEA 450 Scenario.




Figure 1. Global emissions scenarios. (a) fossil CO₂ and (b) all GHGs in CO₂-eq, with gases weighed by their 100-year GWP.

In our scenarios we do not explicitly consider the “negative emissions technologies” to remove CO₂ from the atmosphere, such as an industrial process of direct air capture (e.g., Zeman, 2007), or using biomass for energy and capturing and storing the carbon when the biomass is burned (e.g., Azar et al., 2010). Economics and technical aspects of such options are highly uncertain (House et al., 2011). In addition, there are issues of biomass availability and storage capacity to store carbon at a required scale (Azar et al., 2010). We also do not consider geo-engineering options, such as, for example, solar radiation management (e.g., van Vuuren and Stehfest, 2013), because of the high risks, some of which are known, such as ozone depletion from the introduction of geoengineering aerosol into the stratosphere, but many are still unknown.

3. RESULTING ATMOSPHERIC CONCENTRATIONS

We simulate the atmospheric concentration (**Figure 2**) and climate implications of these scenarios using the MIT IGSM with median setting for earth system response in terms of climate sensitivity, aerosol forcing, and ocean response. All of these scenarios have concentrations levels remaining above 350 ppm through to the end of the century, even when we immediately drop all emissions to zero (**Figure 2a**). In that scenario CO₂ concentrations fall to about 360 ppm by 2100. The gradual decline in concentrations in scenarios where CO₂ emissions drop to zero reflects primarily the initial imbalance between levels of CO₂ in the atmosphere and in the ocean.

The decline in concentrations is relatively rapid in early years as CO₂ levels in the mixed layer of the ocean are more strongly out of balance with levels in the atmosphere. As the ocean's mixed layer comes into balance with the atmosphere, ocean uptake of CO₂ slows. Mixing into the deep ocean is much slower and will continue for hundreds to thousands of years. For example, in the scenario where emissions drop immediately to zero the CO₂ concentration falls by about 10 ppm in the first decade after emissions are cut; this rate of decline falls to only about 1 ppm over the final decade of the century. Eventually this scenario would result in concentrations of 350 ppm of CO₂ or lower, but it would take another 100 years or more.




Figure 2. (a) CO₂ and (b) CO₂-equivalent concentrations. CO₂-eq determined as the CO₂ concentration that would produce the same radiative forcing as the mix of IPCC gases in the atmosphere in that year.

In the Alternative Abatement Scenario proposed as a candidate for achieving a 350 ppm concentration level, we see that instead of falling, concentrations continue to rise from present reaching about 432 ppm by 2035, and remain above 410 ppm through 2100 even with emissions continuing to decline to zero. Overall, these results illustrate Sterman's observation (Sterman, 2008) that the heuristics used in these stock-flow problems are often inaccurate—with the type of error we see here being more the rule than the exception. While the Alternative Abatement Scenario obviously fails to reduce levels to near the 350 ppm goal it was intended to achieve, concentrations are much lower than the Copenhagen Scenario, which reaches about 470 ppm by 2035, and continues rising to over 750 ppm. With emissions dropping to zero in 2035, concentrations only drift down to about 410 ppm by 2100. The IEA 450 Scenario is mostly

successful in keeping CO₂ concentrations in the 450 ppm range, exceeding it by about 15 ppm at the concentration peak, if emissions continue downward toward zero.

Concentrations of all greenhouse gases in CO₂ equivalent behave somewhat differently (**Figure 2b**). First, the Alternative Abatement Scenario reaches over 540 ppm by 2035 but is close to 600 ppm in the Copenhagen Scenario, with the IEA 450 Scenario reaching about 570 ppm in 2035. Concentrations continue to rise for some time, with the Alternative Abatement Scenario and the IEA 450 Scenario topping out at about 550 and 615 ppm respectively. The Copenhagen Scenario continues to rise indefinitely and reaches 1150 ppm by 2100. By 2100 the CO₂-eq concentrations for the IEA 450 Scenario and the Alternative Abatement Scenario are similar to their CO₂ only concentrations, as they are for the Natural Only after 2035 Scenario and Natural Only after 2015 Scenario. In fact, the Natural Only after 2015 Scenario actually achieves a 350 ppm CO₂-eq level by 2100, below the CO₂ alone concentration of 350 ppm.

Why does this occur? The answer lies in short-lived substances, primarily methane. Since methane's lifetime is relatively short—12 to 15 years by most estimates—once emissions drop the concentration level will drop much faster than for longer-lived substances like CO₂. The concentration level in fact drops below preindustrial levels and so becomes a negative contribution to radiative forcing, given that the preindustrial level is used as the benchmark to compute the human contribution to forcing. Concentrations of methane and nitrous oxide are shown in **Figure 3**. Methane concentrations actually fall below preindustrial levels. While preindustrial concentrations are often considered to be unaffected by human activities, there is a body of work documenting likely effects of human activities on GHG concentrations that predate the fossil-fuel combusting, industrial period, mostly ascribing changes to periods of deforestation and reforestation and expansion of paddy rice agriculture (Ruddiman, 2007; Ruddiman et al., 2008). Our representation of anthropogenic versus natural emissions in our IGSM support at least the conclusion that preindustrial methane emissions include a human contribution because that level is not supported only by natural emissions.




Figure 3. (a) CH₄ and (b) N₂O concentrations. Preindustrial concentrations, circa 1750, are represented by the horizontal lines at around 700 ppb and 270 ppb, respectively.

4. RADIATIVE FORCING AND TEMPERATURE EFFECTS

We turn now to the radiative forcing (**Figure 4**) and temperature effects (**Figure 5**) of these five scenarios. In the alternative scenarios we see that radiative forcing is above 8.5 W/m^2 in the Copenhagen Scenario and between 1.9 and 3.6 W/m^2 in stringent scenarios. The mean surface temperature reaches more than $5 \text{ }^\circ\text{C}$ above preindustrial by 2100 in the Copenhagen Scenario. All stringent scenarios keep the temperature increase in 2100 to less than $2 \text{ }^\circ\text{C}$ warmer than the preindustrial mean surface temperature, but they peak before declining. Also note that the temperature keeps warming for about a decade after emissions fall to zero and concentrations peak. This reflects the inertia of the climate system and the removal of negative aerosol forcing. The ocean has a large heat capacity and it takes a significant amount of time to come into equilibrium with changes in forcing. In other words, at any given time the climate system is still responding to the forcing changes in previous years. In addition, aerosol lifetime is much shorter than the lifetimes of greenhouse gases, and when emissions are reduced the immediate reduction in cooling from aerosols leads to an increase in total radiative forcing at that time. With falling concentrations we begin to see a decrease in radiative forcing and after about a decade temperatures begin to fall. In the case where we immediately drop emissions to zero the temperature falls by about $1 \text{ }^\circ\text{C}$ by 2100 from its peak, and remains about half of a degree above preindustrial levels. In the other cases where emissions drop to zero, the temperature falls back to a level between about 1.2 to $2 \text{ }^\circ\text{C}$ above preindustrial. In the Copenhagen Scenario where

emissions continue to rise, temperature continues to rise and shows no sign of slowing. This obviously reflects the fact that concentrations continue to rise.




Figure 4. Radiative forcing. (a) aerosol forcing and (b) total forcing from all GHG gases and aerosols.




Figure 5. Mean surface air temperature for five scenarios. OBS—observations from 1850 to 2010; blue line from 1850 to 2010 represents MIT IGSM reproduction of historic temperature.

5. CONCLUSIONS

As noted at the outset, some have called for CO₂ concentration targets well below the nearly 400 ppm level we have already reached. We have shown here that even if anthropogenic emissions were to drop immediately to zero, levels would still remain above 350 ppm through at least the end of the century. Since CO₂ is not destroyed but rather reallocated among reservoirs—the oceans, atmosphere, and vegetation—our addition to these pools through the combustion of fossil fuels results in, for practical purposes, a permanent increase in the CO₂ that is cycling among these pools. That said, equilibrium with the deep ocean is a very slow process and so over hundreds or thousands of years atmospheric concentrations would continue to slowly decline if we were to reduce emissions to zero. Of course, it is not possible to transform our energy and industrial systems overnight—extremely aggressive policies might imagine such a transformation over 50 years. Of course, we have not considered negative CO₂ emissions technologies, from the simple, such as planting more trees, to the more complex and costly, such as using biomass for energy and capturing and storing carbon when biomass is burned. Somewhat surprisingly we find a 350 CO₂-eq target possibly within range but this depends on eliminating all human influence on methane (and nitrous oxide). Rice agriculture is the biggest human contributor to methane and nitrogen fertilizer use to nitrous oxide. Thus, to achieve these levels requires not only transformation of the world’s energy sector but also its agricultural sector.

Acknowledgments

The MIT Integrated Global System Model (IGSM) and its economic component used in the analysis, the MIT Emissions Prediction and Policy Analysis (EPPA) model, are supported by a consortium of government, industry, and foundation sponsors of the MIT Joint Program on the Science and Policy of Global Change, including U.S. Department of Energy, Office of Science (DE-FG02-94ER61937). (For a complete list of sponsors, see: <http://globalchange.mit.edu/sponsors/all>).

6. REFERENCES

- Azar C., K. Lindgren, M. Obersteiner, K. Riahi, D. Vuuren, K. Elzen, K. Möllersten, and E. Larson, 2010: The feasibility of low CO₂ concentration targets and the role of bio-energy with carbon capture and storage (BECCS). *Climatic Change*, 100: 195–202.
- de Frondeville, B., 2012: Personal communication, April (bertrand@defrondeville.com).
- Gillett, N. P., V. Arora, K. Zickfeld, S. Marshall, and W. Merryfield, W, 2011: Ongoing climate change following a complete cessation of carbon dioxide emissions. *Nature Geoscience*, 4: 83–87.
- Hansen, J., M. Sato, P. Kharecha, D. Beerling, R. Berner, V. Masson-Delmotte, M. Pagani, M. Raymo, D. Royer, and J. Zachos, 2008: Target atmospheric CO₂: Where should humanity aim. *Open Atmos. Sci. J.*, 2: 217–223.
- House K., A. Baslig, M. Ranjan, E. van Nierop, J. Wilcox, and H. Herzog, 2011: Economic and energetic analysis of capturing CO₂ from ambient air. *Proc Natl Acad Sci USA* 108:20428–20433.

- International Energy Agency, 2011: *World Energy Outlook*, OECD/IEA.
- Matthews, H.D., and K. Caldeira, 2008: Stabilizing climate requires near-zero emissions. *Geophys. Res. Lett.* 35: L04705.
- Matthews, H.D., and K. Zickfeld, 2012: Climate response to zeroed emissions of greenhouse gases and aerosols. *Nature Climate Change*, 2, 338–341
- Paltsev, S., S. Dutkiewicz, V. Ekstrom, C. Forest, A. Gurgel, J. Huang, V. Karplus, E. Monier, J. Reilly, J. Scott, A. Slinn, T. Smith-Grieco, and A. Sokolov, 2012: 2012 Energy and Climate Outlook. MITJPSG, *Special Report* (<http://globalchange.mit.edu/research/publications/other/special/2012Outlook>).
- Prinn, R., S. Paltsev, A. Sokolov, M. Sarofim, J. Reilly, and H. Jacoby, 2011: Scenarios with MIT Integrated Global System Model: Significant global warming regardless of different approaches. *Clim. Change*, 104(3-4): 515–537.
- Ruddiman, W., 2007: The early anthropogenic hypothesis: Challenges and responses. *Reviews of Geophysics*, 45: RG4001.
- Ruddiman, W., Z. Guo, X. Zhou, H. Wu, and Y. Yu, 2008: Early rice farming and anomalous methane trends. *Quaternary Science Reviews*, 27(13-14): 1291–1295.
- Sokolov, A., C.A. Schlosser, S. Dutkiewicz, S. Paltsev, D. Kicklighter, H. Jacoby, R. Prinn, C. Forest, J. Reilly, C. Wang, B. Felzer, M. Sarofim, J. Scott, P. Stone, J. Melillo, and J. Cohen, 2005: The MIT Integrated Global System Model (IGSM) version 2: model description and baseline evaluation. MIT JPSPGC *Report 124*, July, 40 p. (http://globalchange.mit.edu/files/document/MITJPSPGC_Rpt124.pdf).
- Solomon, S., G. Plattner, R. Knutti, and P. Friedlingstein, 2009: Irreversible climate change due to carbon dioxide emissions. *Proc. Natl Acad. Sci. USA* 106: 1704–1709.
- Sterman, J., 2008: Risk communication on climate: mental models and mass balance. *Science*, 322(5901): 532–533.
- United Nations, 2009: *Copenhagen Accord*, United Nations Framework Convention on Climate Change (http://unfccc.int/meetings/copenhagen_dec_2009/items/5262.php).
- United Nations, 2010: *Cancun Agreements*, United Nations Framework Convention on Climate Change (<http://cancun.unfccc.int/>).
- Van Vuuren D., and E. Stehfest, 2013: What if climate action becomes really urgent? *Climatic Change*, in press.
- Zeman, F., 2007: Energy and material balance of CO₂ capture from ambient air. *Environ Sci Technol*, 41:7558–7563.

REPORT SERIES of the MIT Joint Program on the Science and Policy of Global Change

FOR THE COMPLETE LIST OF JOINT PROGRAM REPORTS:
<http://globalchange.mit.edu/pubs/all-reports.php>

201. **Russia's Natural Gas Export Potential up to 2050**
Paltsev July 2011
202. **Distributional Impacts of Carbon Pricing: A General Equilibrium Approach with Micro-Data for Households**
Rausch et al. July 2011
203. **Global Aerosol Health Impacts: Quantifying Uncertainties**
Selin et al. August 2011
204. **Implementation of a Cloud Radiative Adjustment Method to Change the Climate Sensitivity of CAM3**
Sokolov and Monier September 2011
205. **Quantifying the Likelihood of Regional Climate Change: A Hybridized Approach**
Schlosser et al. October 2011
206. **Process Modeling of Global Soil Nitrous Oxide Emissions**
Saikawa et al. October 2011
207. **The Influence of Shale Gas on U.S. Energy and Environmental Policy**
Jacoby et al. November 2011
208. **Influence of Air Quality Model Resolution on Uncertainty Associated with Health Impacts**
Thompson and Selin December 2011
209. **Characterization of Wind Power Resource in the United States and its Intermittency**
Gunturu and Schlosser December 2011
210. **Potential Direct and Indirect Effects of Global Cellulosic Biofuel Production on Greenhouse Gas Fluxes from Future Land-use Change**
Kicklighter et al. March 2012
211. **Emissions Pricing to Stabilize Global Climate**
Bosetti et al. March 2012
212. **Effects of Nitrogen Limitation on Hydrological Processes in CLM4-CN**
Lee & Felzer March 2012
213. **City-Size Distribution as a Function of Socio-economic Conditions: An Eclectic Approach to Down-scaling Global Population**
Nam & Reilly March 2012
214. **CliCrop: a Crop Water-Stress and Irrigation Demand Model for an Integrated Global Assessment Modeling Approach**
Fant et al. April 2012
215. **The Role of China in Mitigating Climate Change**
Paltsev et al. April 2012
216. **Applying Engineering and Fleet Detail to Represent Passenger Vehicle Transport in a Computable General Equilibrium Model**
Karplus et al. April 2012
217. **Combining a New Vehicle Fuel Economy Standard with a Cap-and-Trade Policy: Energy and Economic Impact in the United States**
Karplus et al. April 2012
218. **Permafrost, Lakes, and Climate-Warming Methane Feedback: What is the Worst We Can Expect?**
Gao et al. May 2012
219. **Valuing Climate Impacts in Integrated Assessment Models: The MIT IGSM**
Reilly et al. May 2012
220. **Leakage from Sub-national Climate Initiatives: The Case of California**
Caron et al. May 2012
221. **Green Growth and the Efficient Use of Natural Resources**
Reilly June 2012
222. **Modeling Water Withdrawal and Consumption for Electricity Generation in the United States**
Strzepek et al. June 2012
223. **An Integrated Assessment Framework for Uncertainty Studies in Global and Regional Climate Change: The MIT IGSM**
Monier et al. June 2012
224. **Cap-and-Trade Climate Policies with Price-Regulated Industries: How Costly are Free Allowances?**
Lanz and Rausch July 2012.
225. **Distributional and Efficiency Impacts of Clean and Renewable Energy Standards for Electricity**
Rausch and Mowers July 2012.
226. **The Economic, Energy, and GHG Emissions Impacts of Proposed 2017–2025 Vehicle Fuel Economy Standards in the United States**
Karplus and Paltsev July 2012
227. **Impacts of Land-Use and Biofuels Policy on Climate: Temperature and Localized Impacts**
Hallgren et al. August 2012
228. **Carbon Tax Revenue and the Budget Deficit: A Win-Win Solution?**
Sebastian Rausch and John Reilly August 2012
229. **CLM-AG: An Agriculture Module for the Community Land Model version 3.5**
Gueneau et al. September 2012
230. **Quantifying Regional Economic Impacts of CO₂ Intensity Targets in China**
Zhang et al. September 2012
231. **The Future Energy and GHG Emissions Impact of Alternative Personal Transportation Pathways in China**
Kishimoto et al. September 2012
232. **Will Economic Restructuring in China Reduce Trade-Embodied CO₂ Emissions?**
Qi et al. October 2012
233. **Climate Co-benefits of Tighter SO₂ and NO_x Regulations in China**
Nam et al. October 2012
234. **Shale Gas Production: Potential versus Actual GHG Emissions**
O'Sullivan and Paltsev November 2012
235. **Non-Nuclear, Low-Carbon, or Both? The Case of Taiwan**
Chen December 2012
236. **Modeling Water Resource Systems under Climate Change: IGSM-WRS**
Strzepek et al. December 2012
237. **Analyzing the Regional Impact of a Fossil Energy Cap in China**
Zhang et al. January 2013
238. **Market Cost of Renewable Jet Fuel Adoption in the United States**
Winchester et al. January 2013
239. **Analysis of U.S. Water Resources under Climate Change**
Blanc et al. February 2013
240. **Protection of Coastal Infrastructure under Rising Flood Risk**
Lickley et al. March 2013
241. **Consumption-Based Adjustment of China's Emissions-Intensity Targets: An Analysis of its Potential Economic Effects**
Springmann et al. March 2013
242. **The Energy and CO₂ Emissions Impact of Renewable Energy Development in China**
Zhang et al. April 2013
243. **Integrated Economic and Climate Projections for Impact Assessment**
Paltsev et al. May 2013
244. **A Framework for Modeling Uncertainty in Regional Climate Change**
Monier et al. May 2013
245. **Climate Change Impacts on Extreme Events in the United States: An Uncertainty Analysis**
Monier and Gao May 2013
246. **Probabilistic Projections of 21st Century Climate Change over Northern Eurasia**
Monier et al. July 2013
247. **What GHG Concentration Targets are Reachable in this Century?**
Paltsev et al. July 2013